

## Basic Trig Identities

- (1)  $\tan \theta = \frac{\sin \theta}{\cos \theta}$        $\csc \theta = \frac{1}{\sin \theta}$        $\sec \theta = \frac{1}{\cos \theta}$        $\cot \theta = \frac{1}{\tan \theta} = \frac{\cos \theta}{\sin \theta}$
- (2)  $\sin(-\theta) = -\sin \theta$        $\cos(-\theta) = \cos \theta$
- (3)  $\sin(\theta + 2\pi) = \sin \theta$        $\cos(\theta + 2\pi) = \cos \theta$   
 $\sin(\theta + \pi) = -\sin \theta$        $\cos(\theta + \pi) = -\cos \theta$   
 $\sin(\frac{\pi}{2} - \theta) = \cos \theta$        $\cos(\frac{\pi}{2} - \theta) = \sin \theta$
- (4)  $\sin^2 \theta + \cos^2 \theta = 1$
- (5)  $\sin(2\theta) = 2 \sin \theta \cos \theta$
- (6)  $\cos(2\theta) = \cos^2 \theta - \sin^2 \theta$
- (7)  $\sin(\theta + \varphi) = \sin \theta \cos \varphi + \cos \theta \sin \varphi$   
 $\cos(\theta + \varphi) = \cos \theta \cos \varphi - \sin \theta \sin \varphi$

## More Trig Identities

- (4')  $\tan^2 \theta + 1 = \sec^2 \theta$        $1 + \cot^2 \theta = \csc^2 \theta$
- (5',6')  $\tan(2\theta) = \frac{2 \tan \theta}{1 - 2 \tan^2 \theta}$
- (6')  $\cos(2\theta) = 2 \cos^2 \theta - 1 = 1 - 2 \sin^2 \theta$   
 $\cos^2 \theta = \frac{1 + \cos(2\theta)}{2}$   
 $\sin^2 \theta = \frac{1 - \cos(2\theta)}{2}$   
 $\tan^2 \theta = \frac{1 - \cos(2\theta)}{1 + \cos(2\theta)}$
- (7')  $\sin(\theta - \varphi) = \sin \theta \cos \varphi - \cos \theta \sin \varphi$   
 $\cos(\theta - \varphi) = \cos \theta \cos \varphi + \sin \theta \sin \varphi$   
 $\tan(\theta + \varphi) = \frac{\tan \theta + \tan \varphi}{1 - \tan \theta \tan \varphi}$   
 $\tan(\theta - \varphi) = \frac{\tan \theta - \tan \varphi}{1 + \tan \theta \tan \varphi}$
- (7'')  $\sin \theta \cos \varphi = \frac{1}{2} \{ \sin(\theta + \varphi) + \sin(\theta - \varphi) \}$   
 $\sin \theta \sin \varphi = \frac{1}{2} \{ \cos(\theta - \varphi) - \cos(\theta + \varphi) \}$   
 $\cos \theta \cos \varphi = \frac{1}{2} \{ \cos(\theta + \varphi) + \cos(\theta - \varphi) \}$   
 $\sin \alpha + \sin \beta = 2 \sin \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}$   
 $\sin \alpha - \sin \beta = 2 \cos \frac{\alpha + \beta}{2} \sin \frac{\alpha - \beta}{2}$   
 $\cos \alpha + \cos \beta = 2 \cos \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}$   
 $\cos \alpha - \cos \beta = -2 \sin \frac{\alpha + \beta}{2} \sin \frac{\alpha - \beta}{2}$

The code here is that, for example, the identities in (4') are easily derived from the identity in (4). The identity in (5',6') is easily derived by dividing the identities in (5) and (6).